

DEVELOPMENT OF EATING BEHAVIORS AMONG CHILDREN AND ADOLESCENTS

Gurpreet Kaur¹ , Raminder Kaur²

¹Asstant Professor, Baba Fraid College Bathinda.(Punjab) India

²Asstant Professor, Baba Fraid College Bathinda.(Punjab) India

ABSTRACT

The prevalence of obesity among children is high and is increasing. We know that obesity runs in families, with children of obese parents at greater risk of developing obesity than children of thin parents. Research on genetic factors in obesity has provided us with estimates of the proportion of the variance in a population accounted for by genetic factors. However, this research does not provide information regarding individual development. To design effective preventive interventions, research is needed to delineate how genetics and environmental factors interact in the etiology of childhood obesity. Addressing this question is especially challenging because parents provide both genes and environment for children. An enormous amount of learning about food and eating occurs during the transition from the exclusive milk diet of infancy to the omnivore's diet consumed by early childhood. This early learning is constrained by children's genetic predispositions, which include the unlearned preference for sweet tastes, salty tastes, and the rejection of sour and bitter tastes. Children also are predisposed to reject new foods and to learn associations between foods' flavors and the postingestive consequences of eating. Evidence suggests that children can respond to the energy density of the diet and that although intake at individual meals is erratic, 24-hour energy intake is relatively well regulated. There are individual differences in the regulation of energy intake as early as the preschool period. These individual differences in self-regulation are associated with differences in child-feeding practices and with children's adiposity. This suggests that child-feeding practices have the potential to affect children's energy balance via altering patterns of intake. Initial evidence indicates that imposition of stringent parental controls can potentiate preferences for high-fat, energy-dense foods, limit children's acceptance of a variety of foods, and disrupt children's regulation of

energy intake by altering children's responsiveness to internal cues of hunger and satiety. This can occur when well-intended but concerned parents assume that children need help in determining what, when, and how much to eat and when parents impose child-feeding practices that provide children with few opportunities for self-control. Implications of these findings for preventive interventions are discussed.

Key Words- . BMI, body mass index.

INTRODUCTION

This article addresses behavioral factors that influence food preferences, food intake, and energy regulation in children. Currently, the prevalence of childhood overweight is high and has increased dramatically since the 1970s. This increased prevalence is of concern because overweight children are at increased risk for social stigmatization, adult obesity, and chronic disease. Obesity shows familial aggregation; the risk of obesity among children of two obese parents is much higher than for children in families in which neither parent is obese. Familial aggregation has focused research attention on genetic factors in obesity, but the rapid secular increase in the prevalence of obesity cannot be attributable to genetic factors. The interaction of genes and environment influences phenotypes for intake and expenditure and suggests that a renewed focus on the family environment may provide information about behavioral factors that contribute to familial aggregation of adiposity.

Between 30% and 50% of the variance in adiposity within a population is attributable to genetic differences. However, these heritability estimates describe populations, not individuals, and do not provide information about the ways genetics and environment interact during development to produce childhood obesity. In the case of childhood obesity, the question is especially complex because 1) genes and environments also tend to be correlated—parents typically provide children with both genetics and environment; and 2) genetic factors can include behavioral predispositions that affect food intake and expenditure. A more complete understanding of what may characterize obesigenic environments for children is needed, and this review of behavioral factors that influence intake and expenditure indicates ways the family environment can interact with genetic predispositions to produce patterns of food preferences,

food consumption, and physical activity that can promote childhood obesity in susceptible individuals. Obesity results when intake exceeds expenditure, and both sides of the energy-balance equation must be considered in concert in research on the etiology of childhood obesity. Unfortunately, scant research has adopted a conjoint focus on intake and expenditure; thus, we have little knowledge about how children's food intake and physical activity may interact to influence energy balance. Our knowledge also is limited by the lack of precision of the national databases on children's energy intake and the absence of a national database on children's physical activity. Finally, among adults and children, diets high in fat and low in complex carbohydrates are associated with greater adiposity. However, despite repeated recommendations to reduce fat intake, adults' and children's diets are still too high in fat and too low in complex carbohydrates. Our analysis of behavioral factors that influence children's energy intake focuses on research on factors that influence the development of food preferences and the controls of food intake in children. This research will reveal why it is difficult to reduce the intake of energy-dense foods, especially dietary fat intake, and increase complex carbohydrate intake. These findings indicate that attempts to restrict and control children's eating and weight to prevent obesity may be iatrogenic, producing the very problem they are intended to avoid—fostering the development of problems of energy balance. This research also suggests some strategies that can be incorporated into the design of preventive interventions.

Early Experience, Learning, and Children's Food Preferences

One of the first choices that parents make that shapes a child's experience with food and flavors is the choice to breastfeed or formula-feed. The perception of flavors in milk also is one of the human infant's earliest sensory experiences, and there is support for the idea that this early experience with flavors has an effect on milk intake and on later food acceptance. Mennella and Beauchamp have reported that flavors in breast milk influence infants' consumption; breast milk flavored with garlic or vanilla increased the time attached to the maternal nipple relative to breast milk produced on a bland diet. Although Mennella and Beauchamp's research has not focused on individual differences in infants' responsiveness to flavors in milk, a potentially fruitful area for research would explore whether infants who differ in nutritive sucking style, which Agras and

colleagues have related to infant adiposity, also differ in their response to flavors in milk or in the extent to which flavors modulate their ingestion of milk. Galef and Henderson's research with rat pups has demonstrated that early experience with the flavors of the maternal diet can affect subsequent preference for and intake of solid foods. Because of repeated early experience with flavors of the maternal diet present in the mothers' milk, rat pups learn to prefer their mothers' diet. Certainly the early sensory experience of breastfed infants is radically different from that of the formula-fed infant, but we know nothing about how this affects subsequent dietary patterns. Formula-fed infants have experience with only a single flavor, whereas breastfed infants are exposed to a variety of flavors from the maternal diet that are transmitted to the milk. The long-term effects of breastfeeding versus formula-feeding remain unexplored, but very limited evidence suggests that the varied flavor experience of breastfed infants can facilitate acceptance of solid foods during the weaning period, with breastfed infants showing greater initial acceptance of new foods than formula-fed infants. This finding is consistent with those showing that early experience with a variety of flavors leads to more ready acceptance of new foods later. Whether infants differing in familial risk for adiposity may differ in their early responsiveness to flavors or in the transition to solid foods has not been explored, but such research could yield valuable information about the developmental history of gene–environment interactions. Thus, infant dietary experience is shaped by infant-feeding decisions and dietary patterns of the mother, and it provides the basis for food acceptance and patterns of intake in infancy. During the first years of life, the infant transitions from an exclusive milk diet to a variety of foods. This transition from univore to omnivore is shaped by the infant's innate preference for sweet and salty tastes, by the rejection of sour and bitter tastes, and by the omnivore's predisposition to associate food flavors with the contexts and consequences of eating. Infants and children also are predisposed to be xenophobic and reject new foods. With the exception of sweet and salty foods, acceptance of new foods does not occur instantly; however, after repeated opportunities to consume new foods, liking for new food generally increases, producing increased intake, although 5 to 10 exposures often are required. These findings emphasize the central importance of early experience with foods and food acceptance; children come to like and eat what is

familiar. What is familiar is what is present in the environment. There are some data in support of similar dietary patterns between parents and children. Oliver and colleagues reported that parental eating habits had an effect on the nutrient intakes of young children in that children whose parents ate diets high in saturated fat also ate diets high in saturated fats. Parents tend to have foods in the home that they like and eat, and with repeated opportunities to eat these foods, young children include many of them in their diets. The food environment the parent provides shapes children's preferences and food acceptance patterns, which in turn are linked to children's adiposity; we argue that these effects are mediated primarily by the patterns of preference that children have developed as a result of these exposure patterns. The early exposure that children have to fruits and vegetables and to foods high in energy, sugar, and fat may play an important role in establishing a hierarchy of food preferences and selection. The work of Baranowski's group also has confirmed that food availability and accessibility was positively related to fruit and vegetable preferences and to their consumption by school children. These authors noted that children consumed more fruits and vegetables at schools where more fruits and vegetables were served, and they concluded that the extent to which fruits and vegetables are made available and accessible to children may shape children's liking for and consumption of those foods. However, although availability and accessibility are necessary for acceptance of many foods, the physiologic consequences of eating as well as the social context also play a role.

Food Preferences: Predispositions to Prefer Energy-dense Foods

Although repeated exposure can enhance liking for a wide variety of new foods, other learning mechanisms favor development of preferences for energy-dense foods. There is no evidence that children have an innate, unlearned preference for high-fat or high-energy foods (see Birch for a discussion of this point). Children are predisposed to learn to prefer energy-dense foods over energy-dilute foods by learning to associate the flavors of these foods with the positive physiologic consequences that result from eating energy-dense foods, especially when they are hungry. Although limited, the findings from research with young children are consistent with extensive data in the literature on conditioned preferences for energy-dense foods in animals.

Effects of Television on Children's Food Preferences and Food Selection

Television is a pervasive purveyor of culture, providing children with a wide array of models and messages about eating that can influence children's food preferences and food selection as well as their activity patterns. However, despite the central role of television as a purveyor of American culture, surprisingly few studies have investigated the effect of television on children's food preferences, intake, and adiposity. Although there do not appear to be dramatic changes in watching television by children during the period of increased prevalence of childhood obesity, the largest share of advertisements during children's programming is for food products. In 1987, in a content analysis of food advertisements that were on television during a 12-hour period, Cotugna observed that 80% of advertisements showed foods with low nutritional value, including breakfast cereals high in simple sugars and snack foods high in sugar, fat, and salt. A number of investigations have revealed that children's requests for foods were related to the frequency with which children saw the foods advertised on television. Goldberg et al observed that children who were exposed to advertisements selected more sugared foods than children who had not viewed any advertisements. Thus, repeated exposure to food advertisements for particular types of foods may foster children's preferences for energy-dense, nutrient-poor foods. We know even less about how the depiction of eating, dieting, and exercise patterns may influence children's food intake and activity patterns.

Control to Encourage or Limit Children's Eating

Child-feeding practices that control what and how much children eat also can affect their food preferences. Childfeeding strategies that encourage children to consume a particular food increase children's dislike for that food. Many of the foods that parents encourage children to consume are the fruits and vegetables they would like to see consumed with greater frequency and in greater quantities. Hertzler noted that parents' feedback to children about eating vegetables was associated with children's preferences for fewer vegetables. In carrying out what one clinician described as "good food intentions," parents frequently may limit their children's consumption of "bad" but palatable foods by withholding these foods as punishment. In a recent study assessing parental opinions on the efficacy of using various practices to modify their children's food preferences, 40% of parents spontaneously reported the belief that restricting or

forbidding the consumption of a particular food would decrease their child's preference for that food, a prediction opposite research findings. Contrary to these parental beliefs, restricting children's access to foods does not produce food dislikes for the restricted food; instead, such practices enhance liking and can increase intake. Birch et al found that children's preferences for foods increased after the foods were used as rewards for performing a nonfood-related task; limiting the availability of the reward food promoted children's liking of those foods. Again, these restricted foods are typically “bad” foods, those palatable foods that are high in sugar, fat, and energy and that we would like to see consumed in smaller amounts and on fewer occasions. Restricting children's access to foods actually may promote their overconsumption of those foods. Fisher and Birch found that maternal restriction of children's access to snack foods was related to girls' (but not to boys') consumption of those same foods in an unrestricted setting. Creating dietary habits that include moderation and limited consumption of dietary fat and sugar constitutes a desirable objective of child-feeding. However, the limited evidence on the effects of these practices on children's preferences suggests that restricting children's consumption of “bad” foods and encouraging consumption of “good” foods does not provide a means of achieving these dietary goals. From a developmental perspective, this type of feeding practice may send mixed messages to children, because these same forbidden foods are offered in positive social contexts such as parties, dinners out, and holiday celebrations, and are restricted in others. Finally, restricting access to some foods and encouraging consumption of others may well foster children's categorical thinking about “good” and “bad” foods. In a study that involved focus groups of kindergarten children, Murphy et al concluded that children tended to indicate preference for fatty and sugary foods but that they also identified those foods as being high in fat and “not good for you.”

Parental Control and Children's Self-regulation of Food Intake

In addition to influencing which foods children prefer and select, controlling child-feeding practices may affect children's ability to regulate energy intake and the amount of food consumed. As indicated above, a very early decision parents make is choosing whether to formula-feed or breastfeed their infant. In the United States, formulafed infants show more rapid

growth than breastfed infants, and these differences are of sufficient magnitude that some experts have suggested that different growth norms should be used for breastfed and bottle-fed infants. Fomon speculates that the differences in infant growth may be attributable to differences in intake, with the greater intakes by formula-fed infants resulting from overfeeding as a result of heightened maternal control over the infant's intake. In the case of the formula-fed infant, the mother can see how much formula remains in the bottle and she may be inclined to take control of how much the infant eats, encouraging the infant to finish the bottle. In contrast, the breastfed infant may have more control over the size of the feed because this feedback is not available to the mother. Clearly, maternal feeding practices can influence infant intake. In a series of experiments to investigate whether the energy density of formula influenced infants' intake, Fomon and colleagues varied the energy density of formulas infants were consuming and demonstrated that in the absence of maternal control, infants 6 weeks old adjusted their volume of formula intake, consuming more of energy-dilute than of concentrated formulas, so that total energy intake did not differ across the conditions. Fomon's research suggests that when given the opportunity, infants are capable of being responsive to the energy density of formula and adjusting intake accordingly; however, maternal control can override the infant's regulatory ability. There is some evidence for a parallel relationship between parental control and children's ability to regulate intake during early childhood. Although many children continue to be sensitive to internal signals arising from the energy density of the diet in controlling their food intake, child-feeding practices that encourage or restrict children's consumption of foods may decrease the extent to which children use internal signals of hunger and satiety as a basis for adjusting energy intake. To investigate children's responsiveness to energy density, we have conducted research examining children's intake within individual meals by using two-course preloading protocols as well as by examining children's intake across meals by using 24-hour self-selected energy intakes. To determine whether they could adjust intake in response to the energy density of foods within meals, we had children consume a fixed amount of a first course that varied in energy density, by manipulating fat or carbohydrate content, and looked at effects on children's self-selected food intake in a second course. We predicted that if children were

responsive to the energy density of the first course, they would eat less in the self-selected second course after the high-energy than after the low-energy first course. Across a series of experiments, findings confirmed this prediction providing evidence that children can regulate their intake based on feedback arising from the energy content of foods just consumed. Subsequent research extended our findings to reveal that children's responsiveness to energy density can affect their food intake not only within a meal but also across a series of meals, influencing their 24-hour energy intake. Children showed adjustments in energy intake across successive meals in a manner similar to the adjustments in energy intake observed within meals. These adjustments resulted in relatively consistent coefficients of variation for 24-hour energy intake, which averaged ~10%, compared with the much higher coefficients of variation of ~40% for energy intake at individual meals. Children also adjusted their intake to compensate for reduced energy density of meals produced when we substituted olestra for dietary fat, indicating that at least when energy density is reduced, children are responsive to and adjust intake for the energy dilution. Whether children's ability to adjust their food intake in response to energy density is impaired when diets are very high in fat remains an unanswered question. With respect to whether child-feeding practices can alter children's responsiveness to energy density, our initial findings revealed that children's responsiveness to energy density was diminished when adults used control strategies that focused children on external cues to encourage consumption. We explored children's ability to adjust food intake in response to differences in energy density in two different child-feeding contexts.

Factors Influencing Child-feeding Practices: Gender, Adiposity, and Eating Style

Although the potential of nutritional messages to promote restrictive child-feeding practices exists on a societal level, eating-related issues within families also determine parental use of controlling feeding practices. In discussing children's obesity proneness, Costanzo and Woody assert that parents impose behavioral control in domains of their children's development when 1) the parents have problems regulating their own behavior, 2) the child is perceived to be at risk for developing problematic behavior, and 3) the child demonstrates a lack of self-regulatory behavior. These authors use obesity proneness as an example and contend that the use of

controlling parenting styles impedes children's ability to develop self-regulatory behavior, thereby promoting the problems they attempt to avoid. Subsequent research by Johnson and Birch confirmed the links among child-feeding practices and children's responsiveness to energy density. Children's compensation for energy density in our standard two-coursemeal protocol was used as a measure of individual differences in regulation of energy intake. Parents who reported using a high degree of control over what and how much their children ate had children who showed relatively little evidence of energy regulation; a high degree of parental control was associated with low self-control in children. For girls, energy regulation was related to their adiposity, with thinner girls regulating energy intake more precisely than heavier ones. In addition, for girls, but not for boys, parental control was linked to the girls' adiposity, with parents using more control with heavier girls. Parental control also was linked to the parents' dieting and weight history. Mothers who were more restrained used more control and had daughters (but not so for mothers with sons) who showed little evidence of energy regulation. These early gender differences may be precursors of later gender differences in problems of energy balance, in which the prevalence of eating problems is much higher among women. These findings are limited to middle-class white families; we have no data on whether such relationships exist among other racial, ethnic, or socioeconomic groups. One area for future research is to determine whether racial and socioeconomic differences in the prevalence of obesity in children (reported by Troiano and Flegal) may be mediated in part by racial and socioeconomic differences in parenting practices. Fisher and Birch found that young children's weight for height predicted the degree to which mothers reported restricting their child's intake of snack foods. For girls, the parents' dieting and restrictive eating predicted the level of maternal restriction. The cross-sectional nature of the data cannot address the causal direction of the relationships observed, but the results suggest that controlling child-feeding practices adversely affects children's ability to selfregulate food intake and hence their adiposity, although it is possible that parents of heavier children are more controlling. We are obtaining longitudinal data on this question. The child's eating style also may elicit parental concern and control and influence the extent to which the parent sees the child as at risk for obesity and eating problems.

Stunkard and Kaplan reported that the obese ate at a faster rate than normal individuals. Comparisons of obese and normal children have yielded similar results. Drabman et al reported that obese preschoolers ate at a faster rate, taking more bites, and chewing each bite fewer times. Marston and colleagues reported similar patterns of results for school children.

Emergence of Dieting to Control Weight

By middle childhood, even before the fat deposition that occurs in girls at puberty, gender differences in weight concerns and dieting prevalence apparent in adulthood already are emerging. Societal values equating physical attractiveness and thinness with femininity foster the pervasive trend of dieting among most young women, many of whom do not need to lose weight. By the age of 7, children identify overweight silhouette drawings as being less attractive, having fewer friends, and being less smart than their thin counterparts. Even before puberty, girls report a heightened sense of body dissatisfaction and a desire to be thinner, with dieting behaviors reported among children as young as 9 years of age. Serdula et al reported that 44% of high school girls were dieting. Although the nature of dieting behaviors in young girls is not well characterized, the potential adverse consequences of dieting on physical growth and well-being must be addressed. Early dieting itself may constitute a risk behavior for the development of obesity. Dietary restriction involving cognitive restriction of food intake involves explicit denial of hunger cues and includes stopping eating while still hungry and skipping meals. Dietary restriction has been associated with overeating in adolescents and adults. Westenhoefer and colleagues have suggested that restrained eating may give rise to eating binges through the weakening of satiety cues and the heightened attractiveness of food in restrained eaters. Dieting may be iatrogenic, producing the very outcome it is used to avoid. Dieting, with its self-imposed restriction, has certain parallels to controlling child-feeding practices restricting children's intake. Dietary restraint, or the intent to restrict food intake cognitively, has been associated with a number of adverse psychosocial outcomes. In a study of adolescent girls, Killen found that restrained eaters exhibited high levels of worthlessness, body dissatisfaction, fear of weight gain, and disaffect, and they were heavier and were more physically developed than unrestrained girls. Similarly, Rosen et al found that adolescents' restraint scores were positively associated with

depression, body dissatisfaction, social anxiety, and weight status. This all suggests that attempts designed to reduce the prevalence of childhood obesity must consider the potential costs of approaches that may increase the prevalence of early dietary restriction.

Conclusions

Most experts would agree that obesity results when susceptible individuals are placed in adverse environments. To date, there have been few prospective studies of childhood obesity, and these have not tended to focus on the role of environmental factors and how they interact with genetic predispositions that affect intake and expenditure. There is extensive evidence that children's food intake is shaped by early experience with food and eating, and these findings suggest ways in which parenting practices and the family environment may be promoting obesity. Children's eating is modified by exposure and accessibility of foods; by modeling behavior of peers, siblings, and parents; by the physiologic consequences of ingestion; and by child-feeding practices. In particular, the findings reveal ways children's environments can be obesigenic for susceptible individuals. However, the database is very limited; the research reviewed above has focused primarily on white middle-class children of normal weight. There is little evidence about how factors influencing children's food preferences and intake, especially child-feeding practices, may differ systematically across socioeconomic and racial groups. In addition, little research has been conducted with infants and toddlers, in whom dramatic dietary change is occurring, or with adolescents, in whom most research has focused on eating disorders and, to some extent, on dieting behavior. Both infancy and adolescence are critical periods in the development of obesity, yet we know very little about the development of the controls of food intake during these periods. The research on the developing controls of food intake in children of normal weight allows us to generate hypotheses about ways the child's genetic predispositions may interact with environmental factors to produce childhood obesity. What remains on the research agenda are tests of these hypotheses in prospective, longitudinal studies of childhood obesity, which should investigate the possibility that genetic differences contributing to obesity are manifest in individual differences in feeding and exercise predispositions and should include a joint focus on intake and expenditure (see Kohl and Hobbs). Future research should include

children and their families identified as differing in familial risk for obesity, in whom definitions of risk that include behavioral factors such as parenting practices, parental adiposity, and dieting history should be explored. The etiology of childhood obesity is multifaceted and requires a multidisciplinary approach, including expertise on genetics, environmental and behavioral factors, and on both energy intake and the components of energy expenditure. In cases for which doublelabeled water is used to measure energy expenditure and anthropometric measures are taken over time, the validity of dietary intake measures can be addressed. The conjoint focus on children's intake and physical activity will have the added benefit of allowing us to address issues of measurement error, which is essential to advancing our knowledge base.

REFERENCES

- Troiano RP, Flegal KM. Overweight children and adolescents: description, epidemiology, and demographics. *Pediatrics*. 1998;101:497–504
- Troiano RP, Flegal KM, Kuczmarski RJ, Campbell SM, Johnson CL. Overweight prevalence and trends for children and adolescents: the National Health and Nutrition Surveys, 1963 to 1991. *Arch Pediatr Adolesc Med*. 1995;149:1085–1091
- Garn SM, Clark DC. Trends in fatness and origins of obesity. *Pediatrics*. 1976;57:443–456
- Rosenbaum M, Leibel RL. The physiology of body weight regulation: relevance to the etiology of obesity in children. *Pediatrics*. 1998;101: 525–539
- Bouchard C. Genetics of body fat content. In: Angel A, Anderson H, Bouchard C, Lau D, Leiter L, Mendelson R, eds. *Progress in Obesity Research*. Vol 7. London, UK: John Libbey; 1996:33–41
- Fisher JA, Birch LL. 3–5 Year-old children's fat preferences and fat consumption are related to parental adiposity. *J Am Diet Assoc*. 1995; 95:759–764